



AFTER YOUR TVT

INSTRUCTIONS FOR HOME: An important part of your TVT surgery is the recovery period at home. The following information and suggestions should assist your speedy recovery.

ACTIVITY: No lifting greater than 10 pounds for 6 weeks. You may use the stairs. You may return to work and normal activities, as you feel able as long as you don't lift more than 10 pounds. You may feel tired for several weeks after you return home, plan to get some extra rest.

VAGINA: Place nothing in the vagina for 2 weeks. No tampons, no douches, no intercourse for 2 weeks. There may be some light bleeding or discharge from the vagina as the stitch in the vagina heals. Use sanitary pads as you need to. Contact your surgeon if you are concerned about the bleeding or discharge. Go to emergency if you bleed more than a pad worth of blood in less than an hour.

DIET: Your normal diet.

BATHING: You may shower or bath.

PAIN: You may experience pain / discomfort in the pelvis and vagina. For pain you should start by using an anti-inflammatory such as ibuprofen (also called Advil or Motrin). Use according to the instructions on the bottle. If that isn't sufficient for pain control then add acetaminophen (also called Tylenol) and take that according to the instructions on the bottle. You can take both an anti-inflammatory and acetaminophen at the same time. If your surgeon has given you a prescription medicine for pain then add that to the anti-inflammatory and acetaminophen, as you need to. If your pain is not adequately controlled then contact your surgeon.

BLEEDING: There may be a very small amount of bleeding at the two puncture sites just above your pubic bone. If you have any large amount of bleeding, apply pressure to the area and go to the Emergency Department.

MEDICATIONS: You may have been given a prescription for an antibiotic to prevent a bladder infection. Take that according to the instructions on the bottle and be sure to take them all.

POSSIBLE PROBLEMS: Contact your surgeon or go to the Emergency Department if you have: a feeling that you cannot empty your bladder; redness / bleeding / discharge at the incision sites; fever / chills; or pain / swelling in your legs. Please direct concerns and questions to your surgeon. If you do not understand these instructions please ask your nurse or your surgeon to explain. When at home, should you have any questions or complications, call the Fergus office.

POST-OPERATIVE VISIT: You should see your surgeon in 6 weeks. Contact the Fergus office to book that appointment.

CONTACT INFORMATION:

Dr. Reilly and Dr. Rollings-Scattergood – Fergus office (519) 787-7418

This education tool has been prepared by Nursing Staff of Day Surgery at Groves Memorial Community Hospital in consultation with your surgeon. It is our hope that the instructions will alleviate any concerns or apprehensions you may have during your recovery at home.